



genespin

info@genespin.com - www.genespin.com

DATA SHEET

## >2X Quantitative Master Mix Probes high ROX

2X Quantitative Master Mix, is a 2X premixed, ready-to-use solution (Genespin proprietary formulation) containing Xtra Taq Pol, dNTPs, MgCl<sub>2</sub> and stabilizers optimized for use in real time PCR amplification.

2X Quantitative Master Mix contains ROX for dynamic well factor collection. The addition of ROX has no effect on the PCR reaction efficiency or sensitivity of detection.

<i>cat. no.</i>	<i>amount</i>	<i>note</i>
QSTS-RMMix200	5ml	2X conc.
QSTS-RMMix500	12.5ml	2X conc.
QSTS-RMMix1000	25ml	2X conc.

### FOR RESEARCH USE ONLY

#### SHIPPING

Shipped in green ice.

#### STORAGE

Store at -20C°

avoid freeze/thaw cycles, store dark

Storage at 4 °C for up to 3 months possible.

#### SHELF LIFE

12 months

#### FORM

liquid

#### CONCENTRATION

2X



genespin

info@genespin.com - www.genespin.com

DATA SHEET

## >2X Quantitative Master Mix Probes high ROX

### Assay Set-Up:

Before starting, vortex all components thoroughly to ensure homogeneity.

Prepare a premix for the number of assays you need according to the following protocol:

component	stock conc.	final conc.	20ul reaction	50ul reaction
2X Master Mix	2X	1X	10.0ul	25.0ul
primer for	10uM	100-400 nM	0.2 - 0.8 ul	0.5 - 2.0 ul
primer rev	10uM	100-400 nM	0.2 - 0.8 ul	0.5 - 2.0 ul
DNA Template	-	-	< 500ng	< 500ng
MG Water	-	-	up to 20ul	up to 50ul

### Cycling conditions:

Spin down the tubes/plate briefly to remove bubbles and place them into the cycler.

denaturation	95°C	5 min	1x
denaturation	95°C	10 sec	20-35x
annealing (1) and extension	55-68°C	30 sec	

1)The annealing temperature depends on the melting temperature of the primers used.

2)The elongation time depends on the length of the fragments to be amplified. A time of 1 min/kb is recommended.