

DATA SHEET

> XtraTaq Pol White

cat. no.	amount	note
XSTS-T5XW 250	250 units	5X XtraWhite Buffer
XSTS-T5XW 1000	1000 units	5X XtraWhite Buffer
XSTS-T5XWw 250	250 units	5X XtraWhite Buffer w/o MgCl ₂
XSTS-T5XWw 1000	1000units	5X XtraWhite Buffer w/o MgCl ₂

add nucleotide box

cat. no.	amount	note
XSTSn-T5XW 250	250 units	5X XtraWhite Buffer + dNTPs
XSTSn-T5XW 1000	1000 units	5X XtraWhite Buffer + dNTPs
XSTSn-T5XWw 250	250 units	5X XtraWhite Buffer w/o MgCl ₂ + dNTPs
XSTSn-T5XWw 1000	1000units	5X XtraWhite Buffer w/o MgCl ₂ + dNTPs

An highly processive, recombinant (from E.coli strain), thermostable DNA with a very high efficiency of 5'- 3' polymerase activity and 3'- 5' exonuclease (non-proofreading) activity. Xtra Taq Pol catalyzes the addition of mononucleotide units to the 3'-end of a primer chain, leading to the formation of DNA products that have 3'-overhanging A nucleotides (thus can be used in TA cloning). This enzyme remains functional even after prolonged incubation steps at 95°C. The enzyme is supplied at 5U/µl and comes with 5X XtraWhite new buffer.

5X XtraWhite is a Genespin proprietary formulation, developed for standard and/or high-fidelity amplification of high-GC (>75%) templates. The buffer contains 7.5mM magnesium, PCR enhancers and thickening agents (vortex thoroughly prior to use) and is supplied with 1ml of 6X Orange Loading Dye.

The Orange G dye migrates at the same rate as a duplex DNA fragment of approximately 50 Kbp and does not interfere with DNA migration when it is used as a loading dye for agarose gel electrophoresis.

FOR RESEARCH USE ONLY

UNIT DEFINITION

One unit is defined as the amount of enzyme required to incorporate 10 nanomoles of dNTPs into acid-insoluble material in 30 min at 74°C.

SHIPPING

Shipped in green ice.

STORAGE

Store at -20C°

SHELF LIFE

12 months

FORM

Liquid

CONCENTRATION

5U/ul

> XtraTaq Pol White

component	STS-T5XW 250	STS-T5XWw250	STSn-T5XW250	STSn-T5XWw250
Xtra Taq Polymerase	250 units / 50ul	250 units / 50ul	250 units / 50ul	250 units / 50ul
Standard Buffer	1.5ml 5X Buffer with MgCl ₂	1.5ml 5X Buffer w/o MgCl ₂	1.5ml 5X Buffer with MgCl ₂	1.5ml 5X Buffer w/o MgCl ₂
MgCl ₂	-	500ul / 50mM	-	500ul / 50mM
dNTPs	-	-	100ul / 10mM each	100ul / 10mM each

component	STS-T5XW1000	STS-T5XWw1000	STSn-T5XW1000	STSn-T5XWw1000
Xtra Taq Polymerase	1000 units / 200ul	1000 units / 200ul	1000 units / 200ul	1000 units / 200ul
Standard Buffer	4ml 5X Buffer with MgCl ₂	4ml 5X Buffer w/o MgCl ₂	4ml 5X Buffer with MgCl ₂	4ml 5X Buffer w/o MgCl ₂
MgCl ₂	-	500ul / 50mM	-	500ul / 50mM
dNTPs	-	-	400ul / 10mM each	400ul / 10mM each

Assay Set-Up:

Before starting, vortex all components thoroughly to ensure homogeneity.

Prepare a premix for the number of assays you need according to the following protocol:

component	stock conc.	final conc.	30ul reaction
5X Buffer	5X	1X	6.0ul
dNTPs	10mM each.	200uM	0.6ul
Xtra Taq Polymerase	5U/ul	0.025U/ull	0.2ul
primers	1ug/ul each	50ng/ul each	2ul each
DNA Template	-	-	10-20ng
MG Water	-	-	up to 30ul


Cycling conditions:

Spin down the tubes/plate briefly to remove bubbles and place them into the cycler.

denaturation	95°C	5 min	1x
denaturation	95°C	30 sec	20-35x
annealing (1)	50-68°C	30 sec	
extension (2)	72°C	30sec	

1)The annealing temperature depends on the melting temperature of the primers used.

2)The elongation time depends on the length of the fragments to be amplified. A time of 1 min/kb is recommended.



Competitor

5X XtraWhite

Comparisons of the amplification efficiency of new **5X XtraWhite Buffer** versus Competitor were performed using serial dilutions (1:10, 1:100, 1:1000) of Kiwi DNA and specific primers corresponding to pathogen PSA. The results are shown above. New **5X XtraWhite Buffer** demonstrated excellent specificity and high efficiency in amplifying the pathogen.